

PRINCIPLES FOR OBJECT-LINGUISTIC CONSEQUENCE: FROM LOGICAL TO IRREFLEXIVE

CARLO NICOLAI AND LORENZO ROSSI

ABSTRACT. We discuss the principles for a primitive, object-linguistic notion of consequence proposed by [Beall and Murzi 2013] that yield a version of Curry’s paradox. We propose and study several strategies to weaken these principles and overcome paradox: all these strategies are based on the intuition that the object-linguistic consequence predicate internalizes whichever meta-linguistic notion of consequence we accept in the first place. To these solutions will correspond different conceptions of consequence. In one possible reading of these principles, they give rise to a notion of logical consequence: we study the corresponding theory of validity (and some of its variants) by showing that it is conservative over a wide range of base theories: this result is achieved via a well-behaved form of local interpretation. The theory of logical consequence is based on a restriction of the introduction rule for the consequence predicate. To unrestrictedly maintain this principle, we develop a conception of object-linguistic consequence, which we call *grounded* consequence, that displays a restriction of the structural rule of *reflexivity*. This construction is obtained by generalizing Saul Kripke’s inductive theory of truth (strong Kleene version). Grounded validity will be shown to satisfy several desirable principles for a naïve, self-applicable notion of consequence.

1. INTRODUCTION

Object-linguistic treatments of *consequence* have been extensively investigated in the recent literature: on these approaches, consequence is formalized as a predicate in some first-order language, and principles governing its behaviour are given. These studies are motivated by diverse philosophical aims, ranging from criticisms to paraconsistent theories [Whittle 2004], to deflationism about consequence [Shapiro 2011], to new versions of truth-theoretical paradoxes such as Curry’s paradox [Priest and Routley 1982, Beall and Murzi 2013, Mares and Paoli 2014]. Some of these authors, [Beall and Murzi 2013, Murzi and Shapiro 2015] in particular, also stress the analogy between object-linguistic treatments of consequence, truth, and comprehension, and call for a unified solution of the resulting paradoxes, arguing that substructural approaches are preferable to fully structural ones.¹ In order to conform with the terminology adopted in the literature, we will treat ‘consequence’ and ‘validity’ as synonymous where, crucially, consequence or validity do not necessarily coincide with logical consequence or logical validity.

All these approaches can in fact be seen as investigating different ways in which some conclusion ‘follows from’ some premises. In their recent [2013] paper, Beall and Murzi proposed

Date: November 10, 2016.

¹To emphasize even more the analogy with truth, we note that object-linguistic predicate for consequence is needed for the same purposes that motivate a truth predicate, such as blind ascriptions (‘all the derivations made this morning in the Logic lecture are valid’) or generalizations.

the following *naïve* principles for a primitive validity predicate $\text{Val}(x, y)$:

- (VP) If $\varphi \succ \psi$, then $\succ \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)$
 (VD) $\varphi, \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \succ \psi$

where φ and ψ range over sentences possibly containing Val itself, and $\ulcorner \cdot \urcorner$ is informally understood as a name-forming device. It is not completely clear how to read \succ : [Beall and Murzi 2013] interpret it as an unspecified relation of ‘following from’. They also give (what they see as) an analogue of the disquotation schema for truth:

- (Val-Schema) $\succ \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)$ if and only if $\varphi \succ \psi$

(VP) and (VD) are inconsistent with classical logic, over a sufficiently expressive base theory. Beall and Murzi show this via a variant of Curry’s Paradox, which they call *V-Curry Paradox*. In order to introduce the paradox, let us fix the meaning of \succ as a sequent arrow of a system including (VD) and the axioms of a sufficiently strong syntax theory as initial sequents, and closed under (VP) and the standard logical rules – notably, left contraction and cut. Our syntactic axioms enable us to find a sentence ν that is inter-derivable with $\text{Val}(\ulcorner \nu \urcorner, \ulcorner \perp \urcorner)$, where \perp is some falsity of the base theory. An instance of (VD) is $\nu, \text{Val}(\ulcorner \nu \urcorner, \ulcorner \perp \urcorner) \succ \perp$. By definition of ν and contraction we obtain $\nu \succ \perp$. By applying (VP), we get $\succ \text{Val}(\ulcorner \nu \urcorner, \ulcorner \perp \urcorner)$, hence $\succ \nu$. Therefore $\succ \perp$ follows by cut.

In this paper, we explore different strategies to block the V-Curry and related paradoxes. These strategies fall under a common intuition: starting with some meta-theoretic consequence relations, we internalize them in the object language in ways that capture their fundamental traits. That is, each such strategy corresponds to the acceptance of different principles for Val and to different restrictions of the structural rules. Different solutions to the paradox, then, correspond to different ways of cashing out the idea that the acceptance of a sentence of the form $\text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)$ is ultimately to be explained and justified by the acceptance of some meta-theoretical validity statements, where the acceptance of the latter *does not involve object-linguistic validity principles*.

A natural option to develop this strategy, and block the V-Curry (and related paradoxes), is to apply (VP) *only to logical derivations*. Under this reading, Val becomes a primitive predicate for logical validity. This is the strategy followed by [Ketland 2012]: he axiomatizes Val over Peano Arithmetic (henceforth PA) and proves the consistency of the theory resulting from this restriction of (VP). This option is supported by the fact that (VP) does not preserve *logical* validity. The very possibility of formulating (VP) requires a well-behaved machinery to handle the name-forming device $\ulcorner \cdot \urcorner$ and this machinery does not satisfy uniform substitutivity, violating a basic requirement for logically valid principles.²

However, Ketland’s consistency proof only applies to a restricted category of theories. These theories will be called later reflexive theories. Moreover, Ketland’s strategy is based on the possibility of reducing the primitive logical validity predicate to a provability predicate definable in PA. Besides establishing consistency, such proof-theoretic reductions (such as conservativity and variants of interpretability) also help us in characterizing the notion associated with the

²Suppose in fact that (VP) is logically valid, that there is a logical derivation of $\varphi \succ \psi$, and that f is a function in the language from names of sentences to terms that do not name sentences. Then applying (VP) and uniform substitutivity we conclude that $\text{Val}(f(\ulcorner \varphi \urcorner), f(\ulcorner \psi \urcorner))$ comes out as logically derivable, which is clearly absurd. This is clearly remarked in [Cook 2014].

logical validity predicate. For instance, the reducibility of the truth predicate to the base theory may be used to assess general conceptions of truth such as truth-theoretical deflationism (see [Horsten 2011, Ch. 7]): in the same way, proof-theoretical reductions might be employed to assess deflationary and other general conceptions of logical validity (see [Shapiro 2011]). Therefore, in section 2, we provide more general reduction techniques that, besides yielding a consistency proof, will give us a finer-grained analysis of logical validity axiomatized over a wide range of base theories.³

A noteworthy feature of object-linguistic logical validity is that *iterations* of the validity predicate are not allowed. For if $\varphi \succ \psi$ is logically valid we can conclude $\text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)$ in the theory of logical validity, but from the latter we cannot conclude $\text{Val}(\ulcorner \emptyset \urcorner, \ulcorner \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \urcorner)$. There are, however, different notions of consequence, and notions expressing ‘following from’ more generally, for which iterations are very natural, such as entailment or implication.⁴ A standard option to approximate iterability is resorting to *hierarchies*, namely stratifying the \succ and the validity predicate. For example, [Field (forthcoming)] suggests a hierarchy of validity predicates and sequent arrows, and the following version of Beall and Murzi’s principles, where \succ_β is read as ‘derivability in the theory of validity of level β ’:

$$\begin{array}{ll} (\text{VP})_{\alpha, \alpha+1} & \text{If } \varphi \succ_\alpha \psi, \text{ then } \succ_{\alpha+1} \text{Val}_\alpha(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \\ (\text{VD})_{\alpha, \alpha+1} & \varphi, \text{Val}_\alpha(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \succ_{\alpha+1} \psi \end{array}$$

In §2.4 we will see that this stratified notion of validity may be understood in terms of a hierarchy of *reflection principles* over the starting theory: therefore, not only stratified object-linguistic validity is classically consistent, but it has a natural conceptual analysis in terms of a hierarchy of soundness extensions of the starting theory.

However, like any hierarchical approach, also this proposal suffers from variants of the so-called ‘Nixon-Dean problem’ (see [Kripke 1975], pp. 694-697); consider for example the following case

Speaker A says: ‘the negation of everything I say follows from what Speaker B says’.

Speaker B says: ‘everything I say follows from what Speaker A says’.

As for truth, these cases pose problems for hierarchical and non-self-applicable accounts of consequence.

In §3-4, we develop an approach to object-linguistic validity that overcomes this problem: it blocks paradoxical arguments and, at the same time, avoids restrictions on (VP), recovers a natural version of (VD), and delivers a single and genuinely self-applicable notion of validity. This will be accomplished by an inductive construction that generalizes the one in [Kripke 1975] (strong Kleene version). The fundamental feature of the construction is that the models (for languages with self-applicable validity) it generates do not satisfy the structural rule of *reflexivity*.⁵ The smallest fixed point of our construction yields a notion that we might call *grounded validity*, in that it extends Kripke’s notion of grounded truth (see [Kripke 1975], pp. 694 and 706-707). This is because the meta-theoretical notion of validity that holds in the base

³Ketland’s results will turn out to be special cases of our findings.

⁴See [Quine 1961].

⁵For a quirk of terminology, ‘reflexive’ will be employed in two different senses: (i) as referred to arithmetical systems that prove the consistency of all their finite subsystems, and (ii) as referred to theories that satisfy the structural rule of reflexivity. We apologize for the possible, but unavoidable, confusion.

language determines the extension of the object-linguistic validity predicate. As we argue in §5, grounded validity affords us a natural reading of the naïve validity-theoretical principles.

2. OBJECT-LINGUISTIC VALIDITY AND CLASSICAL LOGIC

One may be tempted to read the rules (VP) and (VD) as characterizing a notion of logical validity or logical consequence. However, it became soon clear that this temptation should be resisted: object-linguistic treatments of logical consequence simply do not give rise to paradox. This is the conclusion reached by [Ketland 2012], [Cook 2014], and echoed by [Field (forthcoming)]. In particular, the former analyze the Curry-like derivation sketched in the introduction and come to the main conclusion that paradox arises when the principles governing this notion (whatever it may be) of primitive consequence are themselves considered to be logically valid.⁶

In the following two subsections we consider strategies to overcome paradox while keeping classical logic. But we do not only aim at (classical) consistency: by considering suitable reduction methods of the theories with primitive validity to the respective base theory or suitable extensions of it, we intend to study the nature of the concept of validity in relation with the inferential resources of the starting base theory. In particular:

- (1) Improving on [Ketland 2012, Cook 2014], we give a uniform method for the conservativity of the theories of logical validity over an arbitrary theory extending Elementary Arithmetic (EA), a very weak arithmetical theory.
- (2) This method will also yield the reducibility of the theory of logical validity to reflexive base theories (e.g. PA as in [Ketland 2012, Cook 2014]) and its local reducibility in finitely axiomatized based theories (e.g. EA itself), in which ‘reduction’ is intended as a well-behaved version of relative interpretability that preserves arithmetical vocabulary.
- (3) Even if logical validity is extended to purely arithmetical consequence, classical logic can consistently be kept by interpreting Val as provability in the base theory. Starting from this observation, we show that the hierarchy of validity predicates hinted at by [Field (forthcoming)] can be naturally interpreted as a hierarchy of local reflection principles for the starting theory.

These formal results suggest in turn that the notions of primitive logical and arithmetical (or syntactic) validity are not only unparadoxical, but that they can be conceptually reduced, either globally or locally, to notions definable in the base theory or extensions in the same language.

2.1. Arithmetical theories and reductions. We now fix some formal details. We work in the language $\mathcal{L} = \{0, S, +, \times, \exp, \leq\}$ of arithmetic. Occurrences of the quantifiers in expressions of the form $(\forall x \leq t) \varphi(x)$ and $(\exists x \leq t) \varphi(x)$ where t does not contain x are called *bounded*. Formulas containing only bounded occurrences of quantifiers are called *elementary formulas*.

⁶It is also clear that the paradox can be derived by avoiding any syntactic/arithmetical assumption. To see this, one can employ a familiar trick due to [Montague 1963]. A finite formulation of the axioms of a finitely axiomatizable T (see below for the choice of T) can be pushed into the instance of the diagonal lemma needed in the derivation of the paradox. Starting with the instance

$$v \leftrightarrow \text{Val}(\ulcorner v \wedge A \urcorner, \perp)$$

of the diagonal lemma, where A is a finite reaxiomatization of T , one easily obtains a version of the paradox that does not rely on the assumption of the logical validity of the underlying arithmetical theory but only of the principles for logical consequence.

All theories considered below will extend Elementary Arithmetic EA (or, equivalently, $\text{I}\Delta_0$ plus the totality of exponentiation).⁷ The class of *elementary functions* \mathcal{E} is obtained by closing the initial functions $\text{zero}(\cdot)$, $\text{suc}(\cdot)$, $+$, \times , 2^x , $P_i^n(x_1, \dots, x_n) = x_i$ with $(1 \leq i \leq n)$, and truncated subtraction $x \dot{-} y$ under the operations of composition and bounded minimalization:

$$H(\vec{x}) = F(G_1(\vec{x}), \dots, G_n(\vec{x})); \quad (\mu t \leq y) P(\vec{x}, t) = \begin{cases} \text{the least } t \leq y \text{ s.t. } P(\vec{x}, t) \\ 0, \text{ if there is no such } t \end{cases}$$

where F, G_1, \dots, G_n are elementary functions and P an elementary predicate. EA has sufficient resources to naturally introduce new relations corresponding to the elementary functions by proving their defining equations. We will therefore freely employ some functional expressions for the relevant elementary operations and relations.

The formalization of the syntax of first-order theories as it is standardly done in, e.g., [Schwichtenberg and Wainer 2012], is carried out without difficulties in EA. In particular, once we show that the standard arithmetization of the syntax can be captured by elementary functions, the fact that EA can Σ_1 -define precisely the elementary functions ensures us that syntactic predicates and notions can be intensionally captured in it.

Unless otherwise specified, throughout this section we fix a Hilbert-style system for first-order logic in which modus ponens is the only rule of inference: $X \vdash \varphi$ then indicates that there is a derivation in this system of φ from sentences in X , logical axioms, and using modus ponens only. Derivations will therefore be sequences of formulas. Also the notion of *relative interpretation* of a theory U presented via an elementary set of axioms into another elementary presented theory V will repeatedly occur: it can be considered as a triple (U, τ, W) , with τ a translation function $\tau: \mathcal{L}_U \rightarrow \mathcal{L}_W$ that maps n -ary relations of \mathcal{L}_U into \mathcal{L}_W -formulas with n free variables, n -ary functions of \mathcal{L}_U into \mathcal{L}_W -formulas with $n + 1$ free variables satisfying the obvious existence and uniqueness conditions, and that relativizes quantifiers to a suitable \mathcal{L}_W -formula $\delta(x)$, the domain of the translation. In addition, (U, τ, W) satisfies

$$\text{if } U \vdash \varphi, \text{ then } W \vdash \bigwedge_{i \in \text{FV}(\varphi)} \delta(x_i) \rightarrow \varphi^\tau$$

for formulas $\varphi \in \mathcal{L}_U$ and $\text{FV}(\varphi)$ the set of free variables of φ .

A relative interpretation preserves the structure of a proof. On many occasions we will employ a more regimented notion of relative interpretation. An interpretation is direct if quantifiers are unrelativized and identity is mapped into identity. Let U and W be such that $\mathcal{L} \subseteq \mathcal{L}_U \cap \mathcal{L}_W$. We say that U is \mathcal{L} -embeddable in W if there is a direct interpretation of U

⁷For more details on EA and elementary functions the reader may consult [Beklemishev 2005, Schwichtenberg and Wainer 2012]. The axioms of EA are the universal closures of the following:

$$\begin{array}{ll} \text{EA1} & 0 \neq Sx \\ \text{EA2} & Sx = Sy \rightarrow x = y \\ \text{EA3} & x + 0 = x \\ \text{EA4} & x + Sy = S(x + y) \\ \text{EA5} & x \times 0 = 0 \end{array} \quad \begin{array}{ll} \text{EA6} & x \times Sy = (x \times y) + x \\ \text{EA7} & \exp(0) = 1 \\ \text{EA8} & \exp(Sx) = \exp(x) + \exp(x) \\ \text{EA9} & x \leq 0 \leftrightarrow x = 0 \\ \text{EA10} & x \leq Sy \leftrightarrow (x \leq y \vee x = Sy) \end{array}$$

In addition, EA features the principle of induction for *elementary formulas* $\varphi(x)$:

$$(\text{Ind}(\Delta_0)) \quad \varphi(0) \wedge \forall x(\varphi(x) \rightarrow \varphi(Sx)) \rightarrow \forall x \varphi(x).$$

into W that leaves the \mathcal{L} -vocabulary unchanged. \mathcal{L} -embedding is a properly stricter notion than relative interpretability.⁸ Finally, U is *locally interpretable* in W if any finite subtheory of U is relatively interpretable in W . The notion of *local \mathcal{L} -embedding* is defined analogously.

2.2. Object-linguistic logical validity. As anticipated, in this subsection we deal with object-linguistic treatments of *logical* validity: that is we focus on theories that will be obtained by restricting (VP) only to purely logical derivations. It is worth remarking here that, since we are employing classical logic, the deduction theorem holds: this makes the presentation of the theories of logical validity smoother.

Definition 2.1. *Let $T \supseteq \text{EA}$ be formulated in $\mathcal{L}_V = \mathcal{L} \cup \{\text{Val}\}$. The theory T^{V_0} extends T with the principles, for all \mathcal{L}_V -sentences φ, ψ :*

$$(VP_0) \quad \text{if } \varphi \vdash \psi, \text{ then } T^{V_0} \vdash \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)$$

$$(VD_0) \quad T^{V_0} \vdash \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \wedge \varphi \rightarrow \psi$$

We refer to the theory $T^{V_0} \upharpoonright$ as the theory obtained from T^{V_0} by allowing only formulas of \mathcal{L} as instances of nonlogical axiom schemata of T .

T^{V_0} results from restricting (VP) to purely logical derivations. However, since conditional introduction will be assumed throughout this section, it is convenient to work with a unary rather than a binary validity predicate.

Definition 2.2 (Primitive logical validity). *Let $T \supseteq \text{EA}$ be formulated in $\mathcal{L}^+ = \mathcal{L} \cup \{V\}$, where V is now intended as a unary predicate. The theory T^V extends T with the principles, for all \mathcal{L}^+ -sentences φ :*

$$(VP_1) \quad \text{if } \vdash \varphi, \text{ then } T^V \vdash V(\ulcorner \varphi \urcorner)$$

$$(VD_1) \quad T^V \vdash V(\ulcorner \varphi \urcorner) \rightarrow \varphi$$

Again we refer to the theory $T^V \upharpoonright$ as the theory obtained from T^V by allowing only formulas of \mathcal{L} as instances of nonlogical axiom schemata of T .

That T^V is no essential modification of T^{V_0} is guaranteed by the following:

Proposition 2.3. *T^V and T^{V_0} are mutually \mathcal{L} -embeddable, and so are $T^V \upharpoonright$ and $T^{V_0} \upharpoonright$.*

Proof. The idea is entirely straightforward. By employing the recursion theorem to translate within Gödel quotes,⁹ we can uniformly replace $\text{Val}(x, y)$ and $V(x)$ with, respectively, $V(\tau_0(x \rightarrow y))$ and $V(\ulcorner 0 = 0 \urcorner, \tau_1(x))$ where τ_0, τ_1 are suitable (elementary) translations that leave the arithmetical vocabulary unchanged and do not relativize quantifiers. The verification that the two translations are in fact \mathcal{L} -embeddings is routine as the following holds, for $i \leq 1$ and φ either in \mathcal{L}^+ or in \mathcal{L}_V :

$$(1) \quad \text{if } \varphi \text{ is provable in pure logic, then so is } \varphi^{\tau_i}$$

⁸ Feferman's theorem on the interpretability of inconsistency (see [Visser forthcoming]) represents a separating example between relative interpretation and \mathcal{L} -embeddings. Let T be ω -consistent: by Feferman's theorem, $T + \neg \text{Con}(T)$ is interpretable in T , but it cannot be \mathcal{L} -embeddable in T because \mathcal{L} -embeddability clearly preserves ω -inconsistency. For a study of how to separate \mathcal{L} -embeddings from stronger notions of equivalence, see [Nicolai forth2.].

⁹We refer to [Halbach 2011, §5.3] for motivation and to [Schwichtenberg and Wainer 2012, §2.6.1] for a proof of the theorem in EA.

□

It is intuitively fairly clear that the derivation of the V-Curry paradox is blocked in T^V . When (VP) is applied in the informal presentation of the paradox on p.2, (VD) has been already employed and therefore (VP) is not applied to a sequent obtained via a purely logical derivation. This indicates a strategy to prove the consistency of T^V , which anticipates some traits of the construction carried out in §3.2. The following is a positive inductive definition of the set S of logical truths of \mathcal{L}^+ :

$$(2) \quad y \in S \leftrightarrow \text{Sent}_{\mathcal{L}^+}(y) \wedge (\text{LAx}(y) \vee \exists x (x \in S \wedge (x \rightarrow y) \in S))$$

where $\text{Sent}_{\mathcal{L}^+}$ and LAx are the set of (codes of) sentences of \mathcal{L}^+ and of logical axioms respectively. A fixed point of this definition is a set X such that (\mathbb{N}, X) models (2) in the sense that X is taken as the extension of S . It is easy to see that the least such fixed point I_V is reached after ω iterations of the operator associated with (2) and that the structure (\mathbb{N}, I_V) is a model of T^V (and $T^V|$).

However, consistency may be seen as a necessary but not sufficient condition for a full characterization of the concept of validity captured by T^V . As we mentioned above, a primitive validity predicate is usually motivated – for instance in [Shapiro 2011, Beall and Murzi 2013, Murzi and Shapiro 2015] – along similar lines as the truth predicate: in both cases we aim at expressing meta-theoretic facts in the object-language. For instance, one might want to prove in T^V that all tautologies are logically valid, or that so are all implications from a finite subsets of the axioms of T^V . A natural question concerns therefore the costs of the extra expressive power given by V with respect to the inferential resources of the base theory. Moreover, it is mathematically interesting to weigh these costs across a wide range of possible syntactic base theories by abstracting away specific conditions related to a particular choice of the base theory. From this point of view, a general study of the properties of the theory of object-linguistic validity such as its \mathcal{L} -embedding in the base theory T , conservativity over T , finite axiomatizability over T , become integral part of the study of this notion of validity.

The analogy with truth cannot be pushed much further; in particular, it would be a mistake to see theories of logical validity as a subspecies of theories of truth. Theories of truth featuring the truth-theoretic version of (VD₁) are usually prone to an asymmetry between the internal theory – i.e. what the theory proves true – and the set of its theorems: they prove the conjunction $\lambda \wedge \neg T^\top \lambda^\top$ for some sentence λ , where T is the truth predicate. In other words, the theory displays the puzzling feature of asserting a sentence while declaring it untrue.¹⁰ The situation in T^V is both similar and radically different. By diagonalization, we can obviously obtain a sentence χ such that

$$(3) \quad T^V \vdash \chi \leftrightarrow \neg V(\ulcorner \chi \urcorner)$$

By (VD₁) and (3), we can derive $\neg V(\ulcorner \chi \urcorner)$ and therefore χ in T^V . However, if V is interpreted as logical validity, it is not only harmless, but even desirable for χ to be derivable in T^V but not logically valid because its derivation crucially involves (VD₁).

Next we show that the primitive notion of consequence given by T^V cannot serve an expressive role of finite re-axiomatization.¹¹

¹⁰This is the standard objection against classical theories such as a version of the well-known theory Kripke-Feferman KF. See [Halbach 2011, Field 2008].

¹¹The same proof applies to $T^V|$.

Lemma 2.4. T^V is not finitely axiomatizable.

Proof. Seeking a contradiction, let T_0 be finite reaxiomatization of T^V such that $T_0 \dashv\vdash T^V$. This entails that we can find a finite subtheory A of T^V such that $A \dashv\vdash T^V$ and in which (VP_1) and (VD_1) can only be applied to sentences of \mathcal{L}^+ containing at most n logical symbols. Let \mathcal{L}_n^+ be this latter set of sentences. Now adapt (2) in the following way, where $\text{Sent}_{\mathcal{L}_n^+}$ is the set of (codes of) sentences of \mathcal{L}^+ containing at most n occurrences of logical symbols:

$$k \in S \leftrightarrow k = \ulcorner \neg(\underbrace{\top \wedge \dots \wedge \top}_{\wedge \text{ applied } n\text{-times}}) \urcorner \vee \left[k \in \text{Sent}_{\mathcal{L}_n^+} \wedge (\text{LAx}(k) \vee \exists m (m \in \text{Sent}_{\mathcal{L}_n^+} \wedge m \in S \wedge (m \rightarrow k) \in S)) \right]$$

Let I_{V^n} be the least fixed point of this inductive definition. Therefore (\mathbb{N}, I_{V^n}) is a model of A but it cannot be a model of T^V . \square

We now move on to the question of the conservativity and \mathcal{L} -embedding of T^V in T . In what follows, we distinguish between *reflexive* and *finitely axiomatizable* extensions of EA: We recall that a theory is *reflexive* if it proves the consistency of any of its finite subtheories: for all natural choices of T , reflexive theories T prove, for finite $S \subset T$ and for all $\varphi \in \mathcal{L}$,

$$\text{Rfn}(S) \quad \text{Pr}_S(\ulcorner \varphi \urcorner) \rightarrow \varphi$$

For reflexive T , the question of the \mathcal{L} -embedding and conservativity of T^V in T is readily obtained: let us define the elementary translation $\alpha: \mathcal{L}^+ \rightarrow \mathcal{L}$:

$$\begin{aligned} (s = t)^\alpha &:= s = t & (\forall t)^\alpha &:= \text{Pr}_\emptyset(\alpha(t)) \\ (\neg\varphi)^\alpha &:= \neg\varphi^\alpha & (\varphi \wedge \psi)^\alpha &:= \varphi^\alpha \wedge \psi^\alpha \\ (\forall x\varphi)^\alpha &:= \forall x\varphi^\alpha \end{aligned}$$

The definition of α again relies on Kleene's recursion theorem: in particular $\alpha(\cdot)$ represents $(\cdot)^\alpha$ in EA.

Lemma 2.5.

- (i) If T is reflexive, then α is an \mathcal{L} -embedding of T^V in T .
- (ii) If T is finitely axiomatizable, α cannot be an interpretation of T in T^V .

Proof. Both proofs are immediate. For (i), one simply notices that T , being reflexive, proves $\text{Pr}_\emptyset(\ulcorner \varphi \urcorner) \rightarrow \varphi$ for all $\varphi \in \mathcal{L}$ by $\text{Rfn}(S)$.

For (ii), if α were an interpretation of T^V in T , by letting A be again a finite axiomatization of T , we would have

$$A \vdash \text{Pr}_A(\perp) \rightarrow \perp,$$

(with $\perp := \ulcorner 0 = 1 \urcorner$), which contradicts Gödel's second incompleteness theorem. \square

Part (i) of the previous lemma obviously entails the interpretability of T^V in T for reflexive T , being \mathcal{L} -embeddings stricter than interpretability. Two further remarks: the interpretation α is a variant of the one contained in [Ketland 2012], which only takes care of external occurrences of \forall without applications of the recursion theorem. Moreover, in Lemma 2.5(i), α is indeed an \mathcal{L} -embedding of T^V in T when the latter is reflexive. This clearly indicates that, in the case of reflexive theories, the notion of validity governed by (VP_1) and (VD_1) can be unequivocally understood as a definable notion of logical validity.

Lemma 2.5(i) also immediately yields the conservativity of T^V over T for reflexive T . But reflexive theories are in many senses very special and they have a peculiar behaviour with respect to interpretability and related notions. For instance, by Orey's compactness theorem,¹² reflexive theories collapse the distinction between local and global interpretability and they have the very convenient feature of proving the reflection principle for pure logic that is, as we have seen, closely related to (VD_1) . It is therefore natural to generalize the picture given by Lemma 2.5 and ask ourselves whether the notion of consequence captured by (VP_1) and (VD_1) can be uniformly characterized also in the case of non-reflexive theories. As we anticipated, we will focus on finitely axiomatized theories, which are provably distinct from reflexive theories due to Gödel's second incompleteness theorem.

We recall that U is *locally interpretable* in V if every finite subtheory of U is interpretable in V . Similarly, U is *locally \mathcal{L} -embeddable* in V if every finite $U_0 \subseteq U$ is \mathcal{L} -embeddable in V . We also recall that T^V is formulated in a Hilbert-style calculus in which modus ponens is the only rule of inference. Proofs in T^V of φ are therefore objects of the form $\mathcal{D} = \langle \varphi_0, \dots, \varphi_{n-1}, \varphi \rangle$ where each element of the sequence is either an axiom of T^V or it has been obtained from previous elements by modus ponens. Also, the code $|\mathcal{D}|$ of \mathcal{D} is the code of $\langle \ulcorner \varphi_0 \urcorner, \dots, \ulcorner \varphi_{n-1} \urcorner \rangle$.

To prove the local \mathcal{L} -embeddability of T^V in T , we need the following, well known fact:

Lemma 2.6 (Σ_1 -completeness). *For Σ_1 -formulas φ of \mathcal{L} and any T extending EA, if $\mathbb{N} \models \varphi$, then $T \vdash \varphi$.*

The informal idea for the proof of the local \mathcal{L} -embeddability of T^V in T is straightforward: we translate only the outermost occurrences of \forall because only one 'layer' of the logical validity predicate matters in logical proofs. This enables us to dispense with uses of more sophisticated devices, such as the recursion theorem, to translate within Gödel corners.

Proposition 2.7. *T^V is locally \mathcal{L} -embeddable in T .*

Proof. Let B be a finite subsystem of T^V . In B , we can safely assume that there are at most m applications of (VP_1) to logical proofs \mathcal{D}_i , $i \leq m$.

Let $|\mathcal{D}_i| \leq n$ for all *logical proofs* \mathcal{D}_i , that is, the code of each such \mathcal{D}_i is smaller or equal than n . By our assumptions on sequence coding (see [Schwichtenberg and Wainer 2012, §2.2]), bounds for (codes of) sequences and their concatenations are given by

$$\underbrace{\langle k, \dots, k \rangle}_{m\text{-times}} \leq (k+1)^{2^m}; \quad s_0 \widehat{\ } s_1 \leq (s_0 + s_1)^{2^{2\text{lh}(s_0)}}.$$

where $\text{lh}(\cdot)$ is the elementary function that outputs the number of the elements of a sequence.

We define an elementary predicate $V_n(x)$ stating that x is proved in predicate logic with a proof whose code is less than n :

$$V_n(x) \text{ :} \leftrightarrow (\exists y \leq n) (\text{Prf}_\emptyset(y, x))$$

Here $\text{Prf}_\emptyset(\cdot, \cdot)$ is elementary and expresses Hilbert-style provability in $\text{PL}(\mathcal{L}^+)$, predicate logic in the language \mathcal{L}^+ . This n is fixed and will be kept so throughout the proof.

¹²The theorem states that if V is reflexive and any finite $U_0 \subseteq U$ is interpretable in V , also U is interpretable in V (see [Hájek and Pudlák 1998, §III]).

We specify the translation \mathfrak{b} : it is important to notice that here we are *not* employing the recursion theorem.

$$\begin{aligned} (s = t)^{\mathfrak{b}} &:= s = t & (\forall x)^{\mathfrak{b}} &:= \forall_n(x) \\ \cdot^{\mathfrak{b}} \text{ commutes with prop. connectives} & & (\forall x\varphi)^{\mathfrak{b}} &:= \forall x\varphi^{\mathfrak{b}} \end{aligned}$$

To verify that \mathfrak{b} is an \mathcal{L} -embedding, we check that (VP_1) and (VD_1) hold modulo the translation. More generally, we show by induction on the length of the derivation in B that, for all \mathcal{L}^+ -sentences φ ,

$$(4) \quad \text{if } B \vdash \varphi, \text{ then } T \vdash \varphi^{\mathfrak{b}}$$

It is clear that we obtain (4) when φ is a logical or arithmetical axiom of B . Let φ be of the form $\forall(\ulcorner\psi\urcorner) \rightarrow \psi$. Obviously we have

$$\text{either } \mathbb{N} \models \forall_n(\ulcorner\psi\urcorner) \text{ or } \mathbb{N} \models \neg\forall_n(\ulcorner\psi\urcorner).$$

If the latter, then we are done by Lemma 2.6 because \forall_n is elementary. If the former disjunct obtains, there is a purely logical derivation \mathcal{D}_j of ψ such that $|\mathcal{D}_j| \leq n$. Then there is a purely logical derivation $\mathcal{D}_j^{\mathfrak{b}}$ of $\psi^{\mathfrak{b}}$ obtained by translating its elements.¹³

For the induction step, we only need to worry about (VP_1) . Now if φ is obtained by an application of (VP_1) , it has the form $\forall(\ulcorner\psi\urcorner)$ for ψ an \mathcal{L}^+ -sentence and there is a purely logical proof \mathcal{D}_i of ψ . By assumption, $|\mathcal{D}_i| \leq n$, therefore $T \vdash \forall_n(\ulcorner\psi\urcorner)$ by Lemma 2.6. \square

Now Proposition 2.7 immediately yields, besides the consistency of T^{\forall} that wasn't seriously doubted, the conservativity of T^{\forall} over T for *any* T extending EA.

Corollary 2.8. T^{\forall} is a conservative extension of T .

Proof. If $T^{\forall} \vdash \varphi$ and $\varphi \in \mathcal{L}$, then already a finite subsystem $B \subset T^{\forall}$ proves φ . By Proposition 2.7, $T \vdash \varphi^{\mathfrak{b}}$. But $\varphi^{\mathfrak{b}}$ is nothing more than φ itself by definition of \mathfrak{b} . \square

It should be noted that the conservativity of $T^{\forall}|$ can be obtained in a straightforward way since any model \mathcal{M} of $T \supseteq \text{EA}$ can be expanded to a model (\mathcal{M}, S) of $T^{\forall}|$ where S is the set specified in the inductive definition (2). It is not clear to us whether this strategy can be adapted to the full T^{\forall} . The strategy employed in Proposition 2.7, however, has the additional advantage of being formalizable with only weak arithmetical assumptions.

Moreover, we obtain another proof of the interpretability of T^{\forall} in T , for T reflexive, by Orey's compactness theorem:

Corollary 2.9. For $T \supseteq \text{EA}$ and reflexive, T^{\forall} is interpretable in T .

As far as the authors know, the question of the global interpretability of T^{\forall} for arbitrary $T \supseteq \text{EA}$ is still open.

¹³By employing the sequence encoding sketched above, one can estimate for the translated proof: $|\mathcal{D}_j^{\mathfrak{b}}| \leq G(n)$, where

$$G(n) = \underbrace{\langle n(n+1)^{2^n}, \dots, n(n+1)^{2^n} \rangle}_{n\text{-times}} = \left(n(n+1)^{2^n} + 1 \right)^{2^{(n(n+1)^{2^n})}}$$

2.3. Extending logical consequence. It seems natural to wonder whether the reduction methods considered in the previous section can be tweaked to satisfy more principles for \forall . As noticed already by Ketland, the \mathcal{L} -embedding α , without essential modifications, gives us a more substantial theory of logical consequence over reflexive theories encompassing principles like

$$(K) \quad \forall(\ulcorner \varphi \rightarrow \psi \urcorner) \wedge \forall(\ulcorner \varphi \urcorner) \rightarrow \forall(\ulcorner \psi \urcorner)$$

$$(5) \quad \neg \forall(\ulcorner \forall(\ulcorner \varphi \urcorner) \urcorner)$$

The consistency of the theory $T^{\forall} + K + 5$ is guaranteed by the following corollary to Lemma 2.5:

Observation 2.10. *The theory $T^{\forall} + K + 5$ is \mathcal{L} -embeddable in and conservative over T for reflexive $T \supseteq \text{EA}$.*

Let's abbreviate $T^{\forall} + K$ by writing $T^{\forall+}$. Can we obtain analogues of Proposition 2.7 and Corollary 2.8 for $T^{\forall+}$ over arbitrary $T \supseteq \text{EA}$? It turns out that we can, by suitably tweaking the proofs given above.¹⁴ The fundamental idea is to modify the bound given in the definition of V_n in Proposition 2.7 to allow for the concatenation of the logical proofs of formulas φ and $\varphi \rightarrow \psi$ of \mathcal{L}^+ when the translations of the antecedent of K are assumed.

Proposition 2.11. *$T^{\forall+}$ is locally \mathcal{L} -embeddable in T .*

Proof. As before, let B be a finite subsystem of $T^{\forall+}$. Again, we fix a standard n as bound for the codes of the finitely many logical proofs \mathcal{D}_i preceding an application of (VP_1) .

Let $C_n(x)$ be equivalent to:

$$(\exists y \leq H(n)) \left(\text{Prf}_{\emptyset}(y, x) \wedge (\forall i \leq \text{lh}(y)) (\text{det}((y)_i, y) \rightarrow \text{tru}(y, i) \leq n \wedge (\exists w \leq n) (\text{Prf}_{\emptyset}(w, (y)_i))) \right)$$

where

$$H(n) = \underbrace{\langle n, \dots, n \rangle}_{n^2\text{-times}} \frown \underbrace{\langle n, \dots, n \rangle}_{n^2\text{-times}} = \left(2(n+1)^{2n^2} \right)^{2^{2n^2}}$$

$\text{det}(x, y)$ is an elementary predicate expressing that x is an ‘only detachable’ member of y , that is the proof only ‘cuts’ x via modus ponens and x is not a proper subformula of any other member of y ; $\text{tru}(x, y)$ is an elementary function that takes the initial subsequence of x with y components and outputs its code.¹⁵ Intuitively, $C_n(x)$ expresses that x has a proof in pure logic that (i) applies modus ponens to assumptions that are themselves logically provable with proofs smaller than n and (ii) in which all subproofs of these assumptions are also smaller than n .

As before, we define the translation \mathfrak{c} that, like \mathfrak{b} , only replaces outer occurrences of \forall in proofs, clearly this time with C_n and not V_n .

The proof now proceeds along similar lines as the proof of Proposition 2.7 except, of course, for the case of K . In particular, we want to show, for an arbitrary $\varphi \in \mathcal{L}^+$,

$$(6) \quad T \vdash C_n(\ulcorner \varphi \urcorner) \wedge C_n(\ulcorner \varphi \rightarrow \psi \urcorner) \rightarrow C_n(\ulcorner \psi \urcorner)$$

¹⁴We adapt to the present setting a more general strategy suggested to the authors by Albert Visser.

¹⁵In the definition of C_n , $(x)_y = z$ is an elementary functional expression corresponding to the projection of the y^{th} -element of x .

As before, if one of $C_n(\ulcorner \varphi \urcorner)$ and $C_n(\ulcorner \varphi \rightarrow \psi \urcorner)$ are not true-in- \mathbb{N} , we obtain the claim by Lemma 2.6. If they are both true, then there are proofs \mathcal{D}_0 and \mathcal{D}_1 in $\text{PL}(\mathcal{L}^+)$ of φ and $\varphi \rightarrow \psi$ respecting the conditions above. Since the codes of the detachable members of both proofs will be smaller than n , and so is the number of their subformulas, we can safely assume that

$$|\mathcal{D}_i| \leq \underbrace{\langle n, \dots, n \rangle}_{n^2\text{-times}}$$

for $i \in \{0, 1\}$ and that there is a proof of ψ in $\text{PL}(\mathcal{L}^+)$ with Gödel number $\leq H(n)$. Therefore $\mathbb{N} \models C_n(\ulcorner \psi \urcorner)$ and $T \vdash C_n(\ulcorner \psi \urcorner)$ by Lemma 2.6. \square

By the same argument given in Corollary 2.8:

Corollary 2.12. T^{V^+} is a conservative extension of T .

Again Orey's compactness theorem gives us:

Corollary 2.13. For T reflexive, T^{V^+} is interpretable in T .

The results just presented improve the picture discussed in [Ketland 2012, Cook 2014] and tell us that in many respects – especially if one focuses on conservativity – primitive logical validity is uniformly reducible to the resources of the base theory for a much wider class of theories than the one considered before. However, we were not able to show the interpretability, let alone the \mathcal{L} -embedding, of T^{V} and T^{V^+} in T . This is, however, not an unexpected difficulty: by Orey's and analogous results, there is no gap between local and global interpretability in the context of reflexive theories such as PA. In the case of finitely axiomatizable theories, by contrast, the relationships between these two notions vary considerably and are usually hard to establish.¹⁶

From the point of view of the theory of logical validity the lesson to learn is apparent: the combination of (VP) and (VD) *cannot* be taken to characterize logical validity, which is unparadoxical and uniformly conservative over base theories that contain just a minimum amount of syntactic reasoning.

2.4. Arithmetical consequence and hierarchies. In the previous two subsections we analyzed primitive logical consequence based on an introduction rule (VP₁) for the primitive validity predicate restricted to purely logical proofs. It turns out that no incisions on classical reasoning are needed even if one liberalizes (VP₁) to arithmetical consequence. We now show that by iterating this idea to the transfinite we obtain a symmetry between the hierarchy of validity predicates suggested by [Beall and Murzi 2013, Field (forthcoming)] and the hierarchy of local reflection principles for a starting theory T . For the sake of determinateness, we assume our starting theory to be EA, although the arguments would proceed in an analogous way for any $T \supseteq \text{EA}$.

To define a hierarchy of primitive notions of validity, we assume a notation (OR, <) for ordinals up to Γ_0 , available in EA,¹⁷ a countable stock of predicates $V_a(x)$ – where a ranges over codes of ordinals $\alpha < \Gamma_0$ – that is, we take a Latin alphabet letter to code the corresponding

¹⁶For a study of the asymmetry between finitely axiomatized and reflexive theories in the context of theories of truth, see [Nicolai forth1.].

¹⁷In particular, OR is an elementary set of ordinal codes and < an elementary relation isomorphic to the usual well-ordering of ordinals $< \Gamma_0$.

ordinal in the Greek alphabet. We let \mathcal{L}_0 to be \mathcal{L} itself and $\mathcal{L}_{\alpha+1}$ is $\mathcal{L}_\alpha \cup \{V_a\}$; \mathcal{L}_λ , for λ limit, contains all V_b for $\beta < \lambda$.

Definition 2.14 (Hierarchical validity). *Let $S^0 := \text{EA}$. For successor ordinals, with $\alpha < \Gamma_0$, $S^{\alpha+1}$ in \mathcal{L}_α is defined as follows:*

- (HS1 $_\alpha$) if $S^\alpha \vdash \varphi$, then $S^{\alpha+1} \vdash V_a(\ulcorner \varphi \urcorner)$;
 (HS2 $_\alpha$) $S^{\alpha+1} \vdash V_a(\ulcorner \varphi \urcorner) \rightarrow \varphi$ for all $\varphi \in \mathcal{L}_\alpha$
 (HL) $\bigcup_{\beta < \lambda} S^\beta$ for λ limit.

We briefly comment on the halting point Γ_0 : it is motivated by the availability of a natural notation system in EA [Schwichtenberg and Wainer 2012]. Variations are obviously possible: notations for more ordinals are possible in EA, although the details will bring us too far from our main concerns here. By contrast, if one wants to stick with ordinals that are provably well-ordered in EA, one would need to stop at ω^3 . We claim that this hierarchy of validity predicates is closely related to the following, well-known hierarchy of local reflection principles over EA, again for ordinals $\alpha, \lambda < \Gamma_0$:

$$R^0 := \text{EA}; \quad R^{\alpha+1} := R^\alpha + \text{Rfn}(R^\alpha); \quad R^\lambda := \bigcup_{\beta < \lambda} R^\beta.$$

where

$$\text{Rfn}(T) := \text{Pr}_T(\ulcorner \varphi \urcorner) \rightarrow \varphi \quad \text{for all } \varphi \in \mathcal{L}_T.$$

The formalization of provability for the theories R^α can be carried out in a standard way once a notation for the ordinals is available.

Proposition 2.15. *For $\alpha < \Gamma_0$, S^α is \mathcal{L} -embeddable in R^α .*

Proof. For each $\alpha < \Gamma_0$, we define a translation $\mathfrak{d}: \mathcal{L}_{<\alpha} \rightarrow \mathcal{L}$ as follows, for all $\beta < \alpha$, and where $\mathcal{L}_{<\alpha} := \bigcup_{\beta < \alpha} \mathcal{L}_\beta$:

$$\begin{aligned} (s = t)^\mathfrak{d} &:= (s = t) & (\neg\varphi)^\mathfrak{d} &:= \neg\varphi^\mathfrak{d} \\ (\varphi \wedge \psi)^\mathfrak{d} &:= \varphi^\mathfrak{d} \wedge \psi^\mathfrak{d} & (\forall x\varphi)^\mathfrak{d} &:= \forall x\varphi^\mathfrak{d} \\ (V_b(x))^\mathfrak{d} &:= \text{Pr}_{R^\beta}(\mathfrak{d}(x)) \end{aligned}$$

Now we argue inductively given that $V^0 = R^0$ and that limit stages are not problematic. For HS2 $_\alpha$, if, in $R^{\alpha+1}$, we have $\text{Pr}_{R^\alpha}(\mathfrak{d}(\ulcorner \varphi \urcorner))$ for a standard φ , we can conclude $\varphi^\mathfrak{d}$ since $\text{Sent}_{\mathcal{L}}(\mathfrak{d}(\ulcorner \varphi \urcorner))$, provably in EA. For HS1 $_\alpha$, we can safely assume that $R^\alpha \vdash \varphi^\mathfrak{d}$. Therefore already in EA, $\text{Pr}_{R^\alpha}(\mathfrak{d}(\ulcorner \varphi \urcorner))$. \square

Proposition 2.15 suggests at least the following two remarks: for the reader interested in the mathematical strength of the theories S^α , by a result of [Beklemishev 1995], these theories will prove no more Π_1 -arithmetical sentences than ω^α iterated consistency progressions over EA. At the philosophical level, the theories S^α embody a notion of arithmetical validity corresponding to a proper extension of arithmetical provability in the starting theory EA stratified along ordinal paths that are meaningful in the starting theory. No incision on classical logic is needed at this stage. However, as pointed out also in [Field (forthcoming)], the formulation of the theories S^α relies on how many ordinals we can code in the starting theory. In order

to read off a notion of validity from this stratified picture there seem to be only two options: either validity is inherently stratified, or there is a specific countable ordinal α such that S^α fulfills the requirements we are willing to ascribe to the notion of validity. Neither of these alternatives, however, is completely satisfactory; for one thing, there is no reason to think that validity should be stratified, unless one is happy to concede that also truth is a stratified notion. Moreover, the countable ordinals that are provably well-founded in arithmetical theories vary considerably, and it is highly implausible that the notion of validity should be tied to these implementation details.

This is not to say, however, that stratified validity lacks importance. Even if it doesn't afford a viable notion of validity, it gives us a tool to generate validities starting from valid inferences in the base language. This picture will be improved in the in the next two sections, where we will turn the hierarchical increase of validities into a positive inductive definition. This technical shift will yield a truly self-applicable notion of validity whose extension and properties are independent of how we represent ordinals. As in the case of truth, this will require restricting classical reasoning.

3. A NEW CONSTRUCTION FOR NAÏVE VALIDITY

In this section we propose a way of transcending the stratified picture of validity that generalizes Kripke's method to provide models for languages with a self-applicable truth predicate (see [Kripke 1975]). We will be mainly concerned with providing a class of models that makes the naïve principles for validity consistent, and not so much with formulating effectively presented theories of validity (as in the previous section).¹⁸ The models are obtained via fixed points of an inductive construction. Similarly to what happens in Kripke's theory of truth, we have only one validity predicate that can be introduced without restrictions, via (VP). At a fixed point, (VP) and all the principles that are accepted in the construction can be iterated arbitrarily, thus internalizing *all* the inferences deriving from validities of the base language.

3.1. Initial sequents and rules of inferences. Since, in the perspective of an unstratified picture of validity, (VP) is not in question, what are we do to with the V-Curry Paradox? We have seen in the introduction that the paradox forces a restriction of contraction, cut, or (VD). The idea of transcending the hierarchical conception of validity via an inductive definition is at odds with (VD), even though it calls for an unrestricted acceptance of contraction and cut.¹⁹ For sure, (VD) looks perfectly fine if we read Val as an unspecified 'following from' and clearly also as the notion of logical validity studied in §2.2 and §2.3, but things are different if we accept (VP) unrestrictedly: in so doing, we take Val-statements to represent a naïve notion of consequence, namely meta-inferences that hold in virtue of logical, base-theoretic, and validity-theoretic principles. However, in the presence of full (VD), this idea translates into the acceptance of sentences that we might not want to accept, such as ν .

¹⁸It is nonetheless possible to develop axiomatic theories that are adequate for these models and therefore avoid any arbitrariness in choosing a natural halting point for progressions of theories of stratified validity.

¹⁹Substructural theories compatible with Beall and Murzi's principles are available in the literature: the *non-contractive* theory in [Zardini 2011] (see also [Caret and Weber 2015]) and the *non-transitive* approach of [Ripley 2012] and [Cobreros, Egré, Ripley, van Rooij 2014] (see also [Tennant 2015]) support both (VP) and (VD), and block the derivation of \perp . Strictly speaking, these are theories of naïve truth that feature a conditional obeying conditional versions of (VP) and (VD), but they can be easily adapted to naïve validity.

In the perspective of transcending the hierarchy of validity predicates, one might think that the problem with (VD) is that it allows us to conclude ψ on the *assumption* that φ and $\text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)$ hold. However, if the validity predicate represents meta-theoretical inferences (possibly nested, due to its iterability), one might want to employ an elimination rule that is based on Val-statements that are actually *accepted*, rather than arbitrarily assumed. The following elimination rule for Val embodies this intuition:

(VDm) $\text{If } \Gamma \succ \varphi \text{ and } \Delta \succ \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner), \text{ then } \Gamma, \Delta \succ \psi$

Adopting (VDm), however, is not sufficient to avoid the V-Curry Paradox in the presence of *reflexivity*:

(Ref) $\varphi \succ \varphi$

In fact, (Ref) and (VDm) together immediately yield (VD). So, a proof of \perp is now easy to obtain via a modification of the V-Curry derivation given in the introduction, using (Ref) and (VDm).

How can we avoid this new path to triviality? Our proposed solution consists in the development of a Kripke-style positive inductive definition that, while restricting (VD) and (Ref), consistently satisfies (VP), (VDm), contraction, cut, and indeed every other classically valid rule of inference.²⁰ More generally, our construction will operate uniform restrictions on initial sequents: this harmonizes with the motivation to restrict (VD) outlined above, since arbitrary initial sequents might contain Val-sentences codifying problematic inferences. Rules of inference, by contrast, are safe: if we can control the sequents that we accept, we can adopt all such rules. A Kripke-style construction along the lines of the one developed here has been hinted at in [Field (forthcoming)]. [Meadows 2014] also develops an inductive construction that recovers all Beall and Murzi's principles. His construction also rejects reflexivity but, unlike ours, it is not closed under contraction and cut.²¹

3.2. The KV-construction. The generalization of Kripke's construction we propose here consists in dealing with *sequents* rather than single sentences. By 'sequent', from now on, we will mean an object of the form $\Gamma \Rightarrow \Delta$, where Γ (the antecedent) and Δ (the consequent) are finite sets of \mathcal{L}_V -sentences.

Let us describe our approach informally. The following definitions formalize the intuition outlined above, by enabling us to:

- start with a set of sequents containing at least the ones the form $\Gamma \Rightarrow \Delta$, $s_0 = t_0$ and $s_1 = t_1, \Gamma \Rightarrow \Delta$, for $s_0 = t_0$ an atomic arithmetical truth, and $s_1 = t_1$ an atomic arithmetical falsity,
- apply the operational and structural rules of inference to them,
- internalize the sequents so obtained within the validity predicate, interpreting Val via the rules of inference for the classical material conditional \supset .

This process can be iterated: we apply again the operational, structural, and Val rules, and so on *ad infinitum*. This process reaches a fixed point, which provides the desired interpretation of Val.

²⁰For the theory of inductive definitions, see [Moschovakis 1974]. Non-reflexive approaches to paradoxes did not receive an extensive attention in the literature: some works on the topic include [Greenough 2001, Schroeder-Heister 2012, Schroeder-Heister 2016, French 2016].

²¹For a strengthening of Meadows' approach, see [Pailos and Tajer forthcoming].

Definition 3.1. Let $S \subseteq \omega$, and define the set S^+ as follows.

$n \in S^+$ if:

- (i) $n \in S$; or
- (ii) n is $\Gamma \Rightarrow s = t, \Delta$, and $\mathbb{N} \models s = t$; or
- (iii) n is $\Gamma, s = t \Rightarrow \Delta$, and $\mathbb{N} \models s \neq t$; or
- (iv) n is $\Gamma \Rightarrow \varphi \wedge \psi, \Delta$, and $\Gamma \Rightarrow \varphi, \Delta \in S$, and $\Gamma \Rightarrow \psi, \Delta \in S$; or
- (v) n is $\Gamma, \varphi \wedge \psi \Rightarrow \Delta$, and $\Gamma, \varphi, \psi \Rightarrow \Delta \in S$; or
- (vi) n is $\Gamma \Rightarrow \varphi \vee \psi, \Delta$, and $\Gamma \Rightarrow \varphi, \psi, \Delta \in S$; or
- (vii) n is $\Gamma, \varphi \vee \psi \Rightarrow \Delta$, and $\Gamma, \varphi \Rightarrow \Delta \in S$, and $\Gamma, \psi \Rightarrow \Delta \in S$; or
- (viii) n is $\Gamma \Rightarrow \forall x \varphi(x), \Delta$, and for all $t \in \text{Cter}_{\mathcal{L}_V}$, $\Gamma \Rightarrow \varphi(t), \Delta \in S$; or
- (ix) n is $\Gamma, \forall x \varphi(x) \Rightarrow \Delta$, and for some $t \in \text{Cter}_{\mathcal{L}_V}$, $\Gamma, \varphi(t) \Rightarrow \Delta \in S$; or
- (x) n is $\Gamma \Rightarrow \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner), \Delta$, and $\Gamma, \varphi \Rightarrow \psi, \Delta \in S$; or
- (xi) n is $\Gamma, \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \Rightarrow \Delta$, and $\Gamma \Rightarrow \varphi, \Delta \in S$, and $\Gamma, \psi \Rightarrow \Delta \in S$.

Let $\zeta(n, S)$ abbreviate items (i)-(xi). We can express this definition with an operator $\Psi : \mathcal{P}(\omega) \mapsto \mathcal{P}(\omega)$ defined as $\Psi(S) := \{n \in \omega \mid \zeta(n, S)\}$. The operator Ψ is *increasing* and *monotone*, namely:

- for every $S \subseteq \omega$, we have that $S \subseteq \Psi(S)$;
- for every $S_1, S_2 \subseteq \omega$, if $S_1 \subseteq S_2$, then $\Psi(S_1) \subseteq \Psi(S_2)$.

For every $S \subseteq \omega$, the set

$$S_\Psi := \bigcup_{\alpha \in \text{Ord}} \Psi^\alpha(S)$$

is a *fixed point* of Ψ , since $\Psi(S_\Psi) = S_\Psi$. S_Ψ is said to be the fixed point of Ψ *generated by* S . Let's denote with l_Ψ the fixed point of Ψ generated by the empty set:

$$\text{l}_\Psi := \bigcup_{\alpha \in \text{Ord}} \Psi^\alpha(\emptyset).$$

l_Ψ is the *least fixed point* of Ψ : for every $S \subseteq \omega$, $\text{l}_\Psi \subseteq S_\Psi$.

In the next section, we prove that l_Ψ can be used to interpret \mathcal{L}_V -sequents in a non-trivial way. We will then investigate the behaviour of the structural rules and of the validity-theoretical principles in fixed points of Ψ .

4. MAIN PROPERTIES OF THE KV-CONSTRUCTION

We start by showing that there are consistent fixed points of Ψ . A fixed point S_Ψ is *consistent* if it does not contain the empty sequent $\emptyset \Rightarrow \emptyset$. Consistency typically avoids triviality: if $\emptyset \Rightarrow \emptyset$ is in a fixed point closed under weakening, then *every* sequent is in that fixed point.²²

Proposition 4.1. l_Ψ is consistent.

²²Consistency is typically defined as the absence of a contradiction, but our definition is equivalent to that. We could introduce a connective \neg , interpreting $\neg\varphi$ as $\text{Val}(\ulcorner \varphi \urcorner, \ulcorner 0 = 1 \urcorner)$, and show that the classical rules for negation hold for the so-defined \neg in l_Ψ . Then, it would be easy to show that $\emptyset \Rightarrow \emptyset \notin \text{l}_\Psi$ if and only if there is no \mathcal{L}_V -sentence φ s.t. $\emptyset \Rightarrow \varphi \wedge \neg\varphi \in \text{l}_\Psi$.

Proof. The proof is by induction on the stages I_{Ψ}^{α} of the construction of I_{Ψ} . The claim is trivial for I_{Ψ}^0 and I_{Ψ}^1 . Assuming the claim for the stage I_{Ψ}^{α} , one simply notices that the stage $I_{\Psi}^{\alpha+1}$ is obtained from I_{Ψ}^{α} by adding to it all sequents resulting from an application of the clauses of Definition 3.1. It is then clear that, if $\emptyset \Rightarrow \emptyset$ is not in I_{Ψ}^{α} , no such clause can introduce it in $I_{\Psi}^{\alpha+1}$. The limit case follows straightforwardly from the successor cases. \square

We first notice that every stage I_{Ψ}^{α} of I_{Ψ} is closed under left and right weakening. By construction, any sequent $\Gamma \Rightarrow \Delta$ in I_{Ψ}^{α} is obtained by applying a series of Ψ -clauses to sequents containing arithmetical truths or falsities, with arbitrary side sentences. Therefore, in order to have $\Gamma, \Gamma' \Rightarrow \Delta, \Delta'$ in I_{Ψ}^{α} , we simply consider the same succession of Ψ -clauses applied to starting sequents with Γ' and Δ' as extra side sentences.

Lemma 4.2 (Weakening). *For every ordinal α , if $\Gamma \Rightarrow \Delta$ is in I_{Ψ}^{α} , then for every $\Gamma', \Delta' \subseteq \text{Sent}_{\mathcal{L}_V}$, the sequent $\Gamma, \Gamma' \Rightarrow \Delta, \Delta'$ is in I_{Ψ}^{α} .*

Also, since we are dealing with finite sets, left and right contraction hold for every sequent in every stage of the construction of I_{Ψ} .

Lemma 4.3 (Contraction). *For every ordinal α , if $\Gamma, \varphi, \varphi \Rightarrow \Delta$ is in I_{Ψ}^{α} , then $\Gamma, \varphi \Rightarrow \Delta$ is in I_{Ψ}^{α} . Similarly, if $\Gamma \Rightarrow \psi, \psi, \Delta$ is in I_{Ψ}^{α} , then $\Gamma \Rightarrow \psi, \Delta$ is in I_{Ψ}^{α} .*

A crucial feature of our construction is that, at any stage, sequents are *grounded* in at least one sentence in their antecedent or consequent.

Lemma 4.4 (Groundedness). *For every ordinal α and every sequent $\Gamma \Rightarrow \Delta$, if $\Gamma \Rightarrow \Delta$ is in I_{Ψ}^{α} , then there is at least one sentence φ in Γ such that $\varphi \Rightarrow \emptyset$ is in I_{Ψ}^{α} , or at least one sentence ψ in Δ such that $\emptyset \Rightarrow \psi$ is in I_{Ψ}^{α} .*

Proof. We reason by induction on the construction of I_{Ψ} . The claim is trivial for I_{Ψ}^0 . For I_{Ψ}^1 , the claim is also immediate since this set contains only sequents with atomic arithmetical truths in the consequent or atomic arithmetical falsities in the antecedent. We now assume the claim up to α , and prove it for $\alpha + 1$. Let $\Gamma \Rightarrow \Delta \in I_{\Psi}^{\alpha+1}$ be obtained by applying one Ψ -clause to sequents in I_{Ψ}^{α} . We consider two cases.

We first deal with the Ψ -clause for introducing \forall on the right: in this case $\Gamma \Rightarrow \Delta$ has the form $\Gamma \Rightarrow \Delta', \forall x \varphi(x)$. The sequents in I_{Ψ}^{α} from which $\Gamma \Rightarrow \Delta', \forall x \varphi(x)$ is obtained, then, have the following form:

$$(7) \quad \Gamma \Rightarrow \Delta', \varphi(t_0), \dots, \Gamma \Rightarrow \Delta', \varphi(t_n), \dots$$

By induction hypothesis, for every $\Gamma \Rightarrow \Delta', \varphi(t_i)$ in (7), there is a ψ_i in Γ such that $\psi_i \Rightarrow \emptyset$ belongs to I_{Ψ}^{α} , or a χ_i in $\Delta', \varphi(t_i)$ such that $\emptyset \Rightarrow \chi_i$ belongs to I_{Ψ}^{α} . If, for some i , ψ_i or χ_i are in Γ or Δ' , we are done. If there is no i such that ψ_i or χ_i are in Γ or Δ' , the induction hypothesis gives us that $\emptyset \Rightarrow \varphi(t_i)$ is in I_{Ψ}^{α} for all i . Therefore, an application of the Ψ -clause (ix) gives us that $\emptyset \Rightarrow \forall x \varphi(x)$ is in $I_{\Psi}^{\alpha+1}$, as desired.

If $\Gamma \Rightarrow \Delta$ is obtained via the Ψ -clause (xi) of Definition 3.1, then it has the form $\Gamma', \text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner) \Rightarrow \Delta$ and I_{Ψ}^{α} contains $\Gamma' \Rightarrow \varphi_0, \Delta$ and $\Gamma', \varphi_1 \Rightarrow \Delta$. If the induction hypothesis gives us sequents $\psi \Rightarrow \emptyset$ or $\emptyset \Rightarrow \chi$ where ψ or χ are in Γ' or Δ respectively, we are done. In the only other case, the induction hypothesis gives us that $\varphi_1 \Rightarrow \emptyset$ and $\emptyset \Rightarrow \varphi_0$ are in I_{Ψ}^{α} . By the Ψ -clause (xi) of Definition 3.1, then, we get $\text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner) \Rightarrow \emptyset$ in $I_{\Psi}^{\alpha+1}$. \square

To prove the closure of the stages of the construction of l_Ψ under cut, we need the following inversion lemma.

Lemma 4.5 (Inversion). *For every ordinal α , the following holds:*

- (i) If $\Gamma \Rightarrow \varphi \wedge \psi, \Delta$ is in l_Ψ^α , then $\Gamma \Rightarrow \varphi, \Delta$ is in l_Ψ^α and $\Gamma \Rightarrow \psi, \Delta$ is in l_Ψ^α .
- (ii) If $\Gamma, \varphi \wedge \psi \Rightarrow \Delta$ is in l_Ψ^α , then $\Gamma, \varphi, \psi \Rightarrow \Delta$ is in l_Ψ^α .
- (iii) If $\Gamma \Rightarrow \varphi \vee \psi, \Delta$ is in l_Ψ^α , then $\Gamma \Rightarrow \varphi, \psi, \Delta$ is in l_Ψ^α .
- (iv) If $\Gamma, \varphi \vee \psi \Rightarrow \Delta$ is in l_Ψ^α , then $\Gamma, \varphi \Rightarrow \Delta$ is in l_Ψ^α and $\Gamma, \psi \Rightarrow \Delta$ is in l_Ψ^α .
- (v) If $\Gamma \Rightarrow \forall x\varphi(x), \Delta$ is in l_Ψ^α , then for all $t \in \text{Cter}_{\mathcal{L}_\Psi}$: $\Gamma \Rightarrow \varphi(t), \Delta$ is in l_Ψ^α .
- (vi) If $\Gamma, \forall x\varphi(x) \Rightarrow \Delta$ is in l_Ψ^α , then for some $t \in \text{Cter}_{\mathcal{L}_\Psi}$: $\Gamma, \varphi(t) \Rightarrow \Delta$ is in l_Ψ^α .
- (vii) If $\Gamma \Rightarrow \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner), \Delta$ is in l_Ψ^α , then $\Gamma, \varphi \Rightarrow \psi, \Delta$ is in l_Ψ^α .
- (viii) If $\Gamma, \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \Rightarrow \Delta$ is in l_Ψ^α , then $\Gamma \Rightarrow \varphi, \Delta$ is in l_Ψ^α and $\Gamma, \psi \Rightarrow \Delta$ is in l_Ψ^α .

Proof. We will only consider case (viii). Let $\alpha+1$ be the least ordinal such that $\Gamma, \text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner) \Rightarrow \Delta$ is in $\mathsf{l}_\Psi^{\alpha+1}$. Then either (a) $\Gamma, \text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner) \Rightarrow \Delta$ is obtained by the Ψ -clause (xi) of Definition 3.1 or (b) it is obtained via a different Ψ -clause. In case (a) $\Gamma, \varphi_1 \Rightarrow \Delta$ and $\Gamma \Rightarrow \varphi_0, \Delta$ are in l_Ψ^α and we are done. If (b), there are several sub-cases to consider: we just deal with an application of the Ψ -clause (viii), which yields a sequent of the form $\Gamma, \text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner) \Rightarrow \Delta', \forall x\varphi(x)$, with $\Delta = \Delta', \forall x\varphi(x)$. In this case $\Gamma, \text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner) \Rightarrow \Delta', \varphi(t_i)$ is in l_Ψ^α for every i for some formula $\varphi(x)$. By induction hypothesis we obtain, in l_Ψ^α , sequents of the form

$$(\star) \quad \Gamma, \varphi_1 \Rightarrow \Delta', \varphi(t_i) \qquad (\dagger) \quad \Gamma \Rightarrow \varphi_0, \Delta', \varphi(t_i)$$

for every i . By applying the Ψ -clause (viii) to all sequents of the form (\star) and (\dagger) respectively, we obtain that $\Gamma, \varphi_1 \Rightarrow \Delta', \forall x\varphi(x)$ and $\Gamma \Rightarrow \varphi_0, \Delta', \forall x\varphi(x)$ are in $\mathsf{l}_\Psi^{\alpha+1}$, as desired. \square

Finally, we show that every stage of the construction of l_Ψ is closed under cut.

Proposition 4.6 (Closure under cut). *For every α , if $\Gamma \Rightarrow \Delta, \varphi$ and $\varphi, \Gamma \Rightarrow \Delta$ are in l_Ψ^α , then also $\Gamma \Rightarrow \Delta$ is in l_Ψ^α .*

Proof. The proof is by induction. The case for l_Ψ^0 is trivial. The case for l_Ψ^1 is also immediately obtained since, for $\Gamma \Rightarrow \Delta, \varphi$ and $\varphi, \Gamma \Rightarrow \Delta$ to be in l_Ψ^1 , Γ or Δ have to contain at least one atomic arithmetical falsity or truth respectively.

Let us suppose that, for $\alpha > 0$, $\Gamma \Rightarrow \Delta, \varphi$ and $\varphi, \Gamma \Rightarrow \Delta$ are in $\mathsf{l}_\Psi^{\alpha+1}$. There are three main cases to be considered: (a) in the first, $\Gamma \Rightarrow \Delta, \varphi$ and $\varphi, \Gamma \Rightarrow \Delta$ are obtained by means of a Ψ -clause that introduces φ ; (b) in the second, only one of $\Gamma \Rightarrow \Delta, \varphi$ and $\varphi, \Gamma \Rightarrow \Delta$ is obtained via a Ψ -clause that introduces φ ; (c) in the third, neither $\Gamma \Rightarrow \Delta, \varphi$ nor $\varphi, \Gamma \Rightarrow \Delta$ is obtained via a Ψ -clause that introduces φ .

(a) We consider the case in which φ is of the form $\text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner)$. Therefore the sequents

$$(I) \quad \Gamma, \varphi_0 \Rightarrow \varphi_1, \Delta \qquad (II) \quad \Gamma \Rightarrow \varphi_0, \Delta \qquad (III) \quad \Gamma, \varphi_1 \Rightarrow \Delta$$

are in l_Ψ^α . By the weakening lemma applied to (II), also $\Gamma \Rightarrow \varphi_0, \varphi_1, \Delta$ is in l_Ψ^α . By induction hypothesis, since (I) is in l_Ψ^α , also $\Gamma \Rightarrow \varphi_1, \Delta$ is in l_Ψ^α . Therefore, since (III) is also in l_Ψ^α , $\Gamma \Rightarrow \Delta$ will be in $\mathsf{l}_\Psi^\alpha \subseteq \mathsf{l}_\Psi^{\alpha+1}$ as well, as desired.

- (b) We only consider the case in which φ is $\text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner)$. We assume, moreover, that $\Gamma \Rightarrow \Delta, \text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner)$ is obtained via the Ψ -clause (viii) from sequents in \mathcal{I}_Ψ^α of the form

$$(IV) \quad \Gamma \Rightarrow \Delta', \varphi(t_i), \text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner)$$

for all $i \in \omega$, and that $\text{Val}(\ulcorner \varphi_0 \urcorner, \ulcorner \varphi_1 \urcorner), \Gamma \Rightarrow \Delta$ is obtained via the Ψ -clause (xi) from $\Gamma \Rightarrow \varphi_0, \Delta$ and $\Gamma, \varphi_1 \Rightarrow \Delta$ in \mathcal{I}_Ψ^α . On the one hand, by the inversion lemma applied to all sequents of the form (IV), we obtain that all sequents of the form $\Gamma, \varphi_0 \Rightarrow \Delta', \varphi(t_i), \varphi_1$ are in \mathcal{I}_Ψ^α . By the weakening lemma, since $\Delta = \Delta', \forall x \varphi(x)$,

$$(V) \quad \Gamma, \varphi_0 \Rightarrow \Delta, \varphi(t_i), \varphi_1$$

is in \mathcal{I}_Ψ^α for all $i \in \omega$. On the other, from the fact that $\Gamma \Rightarrow \varphi_0, \Delta$ and $\Gamma, \varphi_1 \Rightarrow \Delta$ are in \mathcal{I}_Ψ^α , the weakening lemma gives us

$$(VI) \quad \Gamma \Rightarrow \varphi(t_i), \varphi_0, \varphi_1, \Delta$$

$$(VII) \quad \Gamma, \varphi_1 \Rightarrow \varphi(t_i), \Delta$$

in \mathcal{I}_Ψ^α . By induction hypothesis, since for all $i \in \omega$ (V) and (VI) are in \mathcal{I}_Ψ^α , also $\Gamma \Rightarrow \varphi(t_i), \varphi_1, \Delta$ is in \mathcal{I}_Ψ^α for all $i \in \omega$. Therefore, since for all $i \in \omega$ (VII) is also in \mathcal{I}_Ψ^α , $\Gamma \Rightarrow \varphi(t_i), \Delta$ will be in \mathcal{I}_Ψ^α for all $i \in \omega$. An application of the Ψ -clause (viii) gives us the desired result.

- (c) We consider the case in which $\Gamma \Rightarrow \Delta, \varphi$ and $\varphi, \Gamma \Rightarrow \Delta$ are obtained by application of the Ψ -clause (viii) to sequents in \mathcal{I}_Ψ^α of the form

$$(VIII) \quad \Gamma \Rightarrow \Delta', \varphi_0(t_i), \varphi$$

$$(IX) \quad \varphi, \Gamma \Rightarrow \Delta'', \varphi_1(t_i)$$

for all $i \in \omega$. By the groundedness lemma applied to all sequents of the form (VIII), we obtain, for each $i \in \omega$, that either there is a sequent $\psi_i \Rightarrow \emptyset$ is in \mathcal{I}_Ψ^α for $\psi_i \in \Gamma$, or there is a sequent $\emptyset \Rightarrow \chi_i$ in \mathcal{I}_Ψ^α with $\chi_i \in \Delta', \varphi_0(t_i), \varphi$. In the former case, we are done by the weakening lemma; in the latter case, if $\chi_i \in \Delta'$ we are also done by weakening, otherwise we reason as follows. If χ_i is $\varphi_0(t_i)$ for all i , an application of the Ψ -clause (viii) gives us that $\emptyset \Rightarrow \forall x \varphi_0(x)$ is in $\mathcal{I}_\Psi^{\alpha+1}$, therefore the claim follows by the weakening lemma. If χ_i is φ for some i , we apply the groundedness lemma to all sequents of the form (IX). By the consistency of \mathcal{I}_Ψ , the induction hypothesis cannot give us $\varphi \Rightarrow \emptyset$ in \mathcal{I}_Ψ^α . In all other possible outcomes of the groundedness lemma applied to all sequents of the form (IX), we reason as we did in the corresponding cases of (VIII). \square

Reflexivity is the only structural rule that does not hold unrestrictedly in \mathcal{I}_Ψ (the proof is a small variant of the V-Curry derivation sketched in the introduction).

Lemma 4.7. \mathcal{I}_Ψ cannot contain all the instances of

$$(Ref) \quad \varphi \Rightarrow \varphi$$

for φ an arbitrary \mathcal{L}_V -sentence.

Lemma 4.7 shows also that dropping reflexivity is ‘best possible’: we have a single structural rule that cannot be consistently accepted.²³ \mathcal{I}_Ψ , however, features a weaker form of reflexivity, which follows immediately from the weakening lemma. For every $\varphi \in \mathcal{L}_V$, we can always find Γ and $\Delta \subseteq \text{Sent}_{\mathcal{L}_V}$ such that $\Gamma, \varphi \Rightarrow \varphi, \Delta \in \mathcal{I}_\Psi$, where Γ and Δ can be taken to be disjoint.

²³[Schroeder-Heister 2016] has recently remarked that a similar restriction of reflexivity in the context of rules for naïve comprehension can avoid paradoxes and, at the same time, make both cut and contraction admissible.

4.1. **Naïve validity in l_Ψ .** Several principles for naïve validity (including the (Val-Schema)⁺ formulated and discussed in [Field (forthcoming)]) are recovered in l_Ψ , in the sense made precise by the following result.

Lemma 4.8. *For every $\varphi, \psi \in \mathcal{L}_V$, and every $\Gamma_0, \Gamma_1, \Delta_0, \Delta_1 \subseteq \text{Sent}_{\mathcal{L}_V}$:*

(VDm) *if $\Gamma_0 \Rightarrow \varphi, \Delta_0$ is in l_Ψ and $\Gamma_1 \Rightarrow \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner), \Delta_1$ is in l_Ψ ,
then $\Gamma_0, \Gamma_1 \Rightarrow \psi, \Delta_0, \Delta_1$ is in l_Ψ .*

(Val-Schema)⁺ *$\Gamma, \varphi \Rightarrow \psi, \Delta$ is in l_Ψ if and only if $\Gamma \Rightarrow \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner), \Delta$ is in l_Ψ .*

Since (Val-Schema)⁺ holds in l_Ψ , it is clear that (VP) and (Val-Schema) are recovered in l_Ψ as well.

(VDm) is not to be understood as a ‘weaker version’ of (VD), since there are theories that validate (VD) but for which (VDm) is too strong and yields triviality. The non-transitive approach of [Ripley 2012] and [Cobreros, Egré, Ripley, van Rooij 2014] is a case in point: adapting the theory developed there to the validity predicate, we see that (VD) holds, while an unrestricted acceptance of (VDm) would trivialize the theory. Essentially, this is because (VDm) incorporates a form of cut, which is clearly inadmissible in a non-transitive approach.²⁴

(VD) is the only validity-theoretical principle that does not hold unrestrictedly in l_Ψ . In fact, if l_Ψ contained all instances of (VD), then it would also contain its instance

$$\nu, \text{Val}(\ulcorner \nu \urcorner, \ulcorner \perp \urcorner) \Rightarrow \perp$$

where ν is the V-Curry sentence $\text{Val}(\ulcorner \nu \urcorner, \ulcorner \perp \urcorner)$. The derivation of the V-Curry paradox (outlined in the introduction) would then give us the sequent $\emptyset \Rightarrow \emptyset$ in l_Ψ , against the consistency of l_Ψ .

In Section 3.1 we suggested that a uniform way to avoid V-Curry-driven triviality consists in restricting our acceptance of initial sequents, avoiding the acceptance of Val-sentences that express inferences that we cannot control. Rules of inference, on the other hand, are safe, since the construction of l_Ψ operates a selection over the sequents that are accepted in the first place. This view sits naturally with a restriction of (Ref) and (VD) (the only schematic inferences amongst the structural rules and the validity-theoretical principles) and an unrestricted acceptance of weakening, contraction, cut, (VP), (VDm), and (Val-Schema)⁺. The results 4.2, 4.3, 4.6, 4.7, and 4.8, establish that l_Ψ realizes the solution to the paradoxes described in Section 3.

It is easy to turn l_Ψ into a proper model of the language \mathcal{L}_V , as it is standardly done with Kripke fixed points. Let the *extension* of the validity predicate generated by l_Ψ , in symbols E_Ψ , be the set of those pairs of \mathcal{L}_V -sentences $\langle \varphi, \psi \rangle$ such that $\emptyset \Rightarrow \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner)$ is in l_Ψ , and the *anti-extension* of the validity predicate generated by l_Ψ , in symbols A_Ψ , be the set of pairs of \mathcal{L}_V -sentences $\langle \varphi, \psi \rangle$ such that $\text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \Rightarrow \emptyset$ is in l_Ψ . The model of \mathcal{L}_V naturally associated with l_Ψ , thus, is $(\mathbb{N}, E_\Psi, A_\Psi)$, and its evaluation clauses can be read off Definition 3.1. In particular, $(\mathbb{N}, E_\Psi, A_\Psi)$ can be associated with a three-valued semantics, say with values

²⁴The choice between (VD) and (VDm) reflects an ongoing debate in the truth-theoretical literature (especially concerning substructural theories of truth) on the ‘correct’ form of *modus ponens* (see [Zardini 2013] and [Ripley 2015]). Two of the main contenders are analogous to (VD) and (VDm):

$$\varphi, \varphi \rightarrow \psi \vdash \psi \quad \text{and} \quad \text{from } \Gamma \vdash \varphi \text{ and } \Delta \vdash \varphi \rightarrow \psi \text{ infer } \Gamma, \Delta \vdash \psi.$$

$\{0, 1/2, 1\}$, where the logical vocabulary is interpreted as in strong Kleene semantics (with Val being treated as the strong Kleene conditional), and where $\Gamma \Rightarrow \Delta$ has a *tolerant-strict* reading, that is whenever every $\varphi \in \Gamma$ have value $1/2$ or 1 , then there is a $\psi \in \Delta$ with value 1 .²⁵

We conclude this subsection by noticing that there are close relations between I_Ψ and the least fixed point of Kripke's construction for truth (strong Kleene version) from [Kripke 1975]. This can be achieved by defining $\neg\varphi$ and $T(\ulcorner\varphi\urcorner)$, respectively, as $\text{Val}(\ulcorner\varphi\urcorner, \ulcorner 0 \urcorner = 1\urcorner)$ and $\text{Val}(\ulcorner 0 \urcorner = 0\urcorner, \ulcorner\varphi\urcorner)$, and by constructing the least Kripke fixed point, in the usual way, for the language \mathcal{L}_V . For every sentence $\varphi \in \mathcal{L}_V$ we will have that:

- (8) if φ is in the extension of T in the least Kripke's fixed point, then $\emptyset \Rightarrow \varphi$ is in I_Ψ ;
 if φ is in the anti-extension of T in the least Kripke's fixed point, then $\varphi \Rightarrow \emptyset$ is in I_Ψ .

(8) indicates that Kripke's least fixed point for truth constitutes a proper fragment of I_Ψ . Clearly, the other direction of (8) does not hold.

4.2. Non-minimal fixed points and extensions. Lemma 4.3 shows that every fixed point of Ψ is closed under contraction. However, this is not so for weakening. The fixed point $\{(\emptyset \Rightarrow \mu)\}_\Psi$, where μ is the sentence $\text{Val}(\ulcorner\mu\urcorner, \ulcorner\mu\urcorner)$, for example, is not closed under weakening (μ is the validity-theoretical analogue of the truth-teller). This shortcoming, however, can be easily fixed. Let Ψ^+ be the monotone operator that results by adding to items (i)-(xi) of Definition 3.1 the following positive elementary clause as a further disjunct:

- (xii) n is $(\Gamma, \Gamma' \Rightarrow \Delta', \Delta)$, and $(\Gamma \Rightarrow \Delta) \in S$.

Let's adapt the notation adopted for Ψ to Ψ^+ . The following result is immediate (the first item follows from Lemma 4.2 and the proof of Proposition 4.6, and the second by an induction on the build-up of S_{Ψ^+}).

Lemma 4.9.

- $I_\Psi = I_{\Psi^+}$.
- For every $S \subseteq \omega$, S_Ψ is consistent if and only if S_{Ψ^+} is consistent.

The properties of I_Ψ transfer to I_{Ψ^+} , and the consistency of a fixed point S_Ψ transfers to the fixed point of Ψ^+ generated by the same set S .²⁶ The operator Ψ^+ , however, guarantees closure under contraction as well as cut.²⁷

Observation 4.10. For every $S \subseteq \omega$, every $\varphi \in \mathcal{L}_V$, and every $\Gamma, \Delta \subseteq \text{Sent}_{\mathcal{L}_V}$:

- (L-Wkn) If $\Gamma \Rightarrow \Delta$ is in S_{Ψ^+} , then $\Gamma, \varphi \Rightarrow \Delta \in S_{\Psi^+}$.
- (R-Wkn) If $\Gamma \Rightarrow \Delta$ is in S_{Ψ^+} , then $\Gamma \Rightarrow \varphi, \Delta$ is in S_{Ψ^+} .
- (L-Ctr) If $\Gamma, \varphi, \varphi \Rightarrow \Delta$ is in S_{Ψ^+} , then $\Gamma, \varphi \Rightarrow \Delta$ is in S_{Ψ^+} .
- (R-Ctr) If $\Gamma \Rightarrow \varphi, \varphi, \Delta$ is in S_{Ψ^+} , then $\Gamma \Rightarrow \varphi, \Delta$ is in S_{Ψ^+} .
- (Cut) If $\Gamma \Rightarrow \varphi, \Delta$ is in S_{Ψ^+} and $\Gamma, \varphi \Rightarrow \Delta$ is in S_{Ψ^+} , then $\Gamma \Rightarrow \Delta$ is in S_{Ψ^+} .

²⁵We are grateful to Paul Egré, Francesco Paoli, and Robert van Rooij for suggesting this point to us. Tolerant-strict consequence is also employed in [Meadows 2014].

²⁶We did not use Ψ^+ in the first place in order to simplify the proofs of the properties of I_Ψ : this has no practical effects, however, since I_Ψ and I_{Ψ^+} are identical.

²⁷This result improves on the structural rules recovered by Meadows in [Meadows 2014].

5. FROM LOGICAL TO GROUNDED CONSEQUENCE

In this paper, we investigated different combinations and modifications of the principles (VP) and (VD), and the corresponding notions of consequence. Starting with a restriction of (VP) corresponding to the notion of logical consequence, we explored ways of keeping the full (VP), thereby restricting (VD), carving out a notion of self-applicable consequence grounded in truths and falsities of the base theory. The key findings of the paper are summarized in Table 1.

Consequence	Key Finding
Logical	(i) Conservativity for all $B \supseteq EA$ (ii) <i>Local</i> \mathcal{L} -embedding for all $B \supseteq EA$
Arithmetical	Analysis of Field’s hierarchy via local reflection principles
Grounded	(i) Consistency of (VP), (VDm), (Val-Schema) ⁺ (ii) Kripke-style theory for Val: development of Ψ (iii) Make sense of Val-principles via I_Ψ and grounded validity

TABLE 1. Key Findings

The rules (VP) and (VD) support a strict reading of \succ as logical derivability, and therefore of Val as the class of the logically valid inferences. On this approach, (VP), namely (VP₁) in the formalism of §2, is restricted to purely logical inferences, while the full (VD) – (VD₁) in §2 – can be consistently kept: from this point of view, we can naturally read (VD) as preservation of truth in logically valid arguments. A sub-theory of our theory of logical validity is the theory given by (VP) and a suitable version (VDm): this theory meets the criterion (suggested by [Field (forthcoming)]) of giving the same reading to both the meta-theoretic notion expressed by \vdash and to the predicate Val. The theory of logical validity is therefore simple and well-behaved: it’s then natural to study in more depth the corresponding notion of object-linguistic validity by comparing it to the inferential resources of the underlying base theory. Corollary 2.8 tells us that logical validity does not have any impact on the underlying syntactic structure: for any theory extending a very weak arithmetical system, (VD) and the restricted version of (VP) do not enable us to prove new syntactic or arithmetical facts. This extends to further principles for logical consequence, like internalized modus ponens, as shown in Corollary 2.12. These results seem to suggest that ‘deflationary’ approaches to logical consequence, such as Shapiro’s [Shapiro 2011], may endorse conservativity requirements for logical validity. This is not to say, however, that the predicate of logical consequence does not play an indispensable expressive role: although the theory of logical validity is uniformly locally interpretable in the base theory, our results *do not* show that it is relatively interpretable in the object theory and therefore, arguably, expressively reducible to it.

As we have seen, the consistency of the theory of logical validity follows from a restriction of (VP) to purely logical derivations: consistency, however, is preserved even if we internalize purely arithmetical derivations. This led us, following [Field (forthcoming)], to investigate a hierarchy of arithmetical consequence predicates. We have shown that every stage in this hierarchy corresponds to a stage of a parallel hierarchy of local reflection principles; as a consequence, the hierarchical notion of validity can be read as iterated arithmetical provability.

The hierarchical approach to consequence suffers from several well-known problems: in particular, there seem to be notions of ‘following from’ that cannot be accounted for in the stratified picture, such as implication. Transcending the hierarchy calls for an unrestricted (VP). We have achieved this via a Kripke-style construction, the KV-construction, in which the unrestricted (VP) is balanced by a rule form of (VD). The least fixed point of the KV-construction, l_Ψ , embodies a notion that, following Kripke’s theory of truth, we might call *grounded validity*, i.e. validity grounded in truths and falsities of the base language, in our case arithmetical truths and falsities.²⁸

The main intuition behind grounded validity is that *first* we have inferences involving non-semantic facts, which we can then combine and iterate to express more complex inferences, crucially including nested occurrences of the consequence relation. At the fixed point l_Ψ , the process of generating more and more acceptable inferences reaches a halt: the set l_Ψ realizes in full the idea of iterating arbitrarily the grounded consequences, and of expressing them in the object-language. To see this, let F_Ψ be the set of \mathcal{L}_V -sentences such that φ is in F_Ψ if and only if $\emptyset \Rightarrow \varphi$ is in l_Ψ . It is immediate to see that, thanks to the fixed-point property of l_Ψ , for all sentences φ, ψ we have that:

$$(9) \quad \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \in F_\Psi \text{ if and only if } \text{Val}(\ulcorner \emptyset \urcorner, \ulcorner \text{Val}(\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \urcorner) \in F_\Psi.$$

(9) follows immediately by Lemma 4.8, that shows that (Val-Schema)⁺ holds unrestrictedly in l_Ψ . In his [Field (forthcoming)], Field rejects (Val-Schema)⁺ on the grounds of an example that, informally, reads thus:

$$(10) \quad \begin{array}{l} \text{It follows from ‘snow is white’ that} \\ \text{it follows from ‘grass is green’ that ‘snow is white’} \end{array}$$

If ‘follows from’ is intended as logical consequence, (10) clearly does not hold. However, (10), and more generally (Val-Schema)⁺, cease to be troubling if we adopt the grounded consequence reading: since snow is white, this truth about the non-semantic vocabulary grounds and justifies the consequence expressed in (10).²⁹

Desirable features such as that expressed in (9) come at a cost: we cannot consistently accept all sequents of the form $\varphi \Rightarrow \varphi$ in l_Ψ . As a consequence, not all sentences of the form $\text{Val}(\ulcorner \varphi \urcorner, \ulcorner \varphi \urcorner)$ are in F_Ψ . However, this restriction is not so implausible in the context of a grounded consequence relation. Some sequents of the form $\varphi \Rightarrow \varphi$, in fact, are to be rejected because they are *ungrounded*. A paradigmatic case of failure of (Ref) involves the V-Curry sentence itself: we do not have $\nu \Rightarrow \nu$ in l_Ψ . In other words, we do not accept that

$$(11) \quad \begin{array}{l} \text{from the fact that (from this sentence it follows that } 0 = 1), \\ \text{it follows that (from this sentence it follows that } 0 = 1). \end{array}$$

If we only accept grounded Val-sentences, we want to *unpack* the ‘it follows’ used in (11), to see from where it derives. Given the ungrounded nature of ν , such unpacking does not lead us to non-semantic inferences, but to an endless, circular regress. Cases such as (11) are the only kind of sequents of the form $\varphi \Rightarrow \varphi$ that are not in l_Ψ , which makes the restriction of reflexivity less

²⁸See [Kripke 1975], p. 694 and p. 701. For an analysis of Kripkean groundedness, see [Yablo 1982]. For arguments for Kripkean grounded truth, see [Leitgeb 2005], [Martin 2011], and [Burgess 2014].

²⁹Grounded validity is clearly distinct from analytical validity, i.e. validity based on analytical truths and falsities. In fact, it is possible to start our construction from non-analytic claims, e.g. about snow being white and grass being green. Thanks to Andreas Fjellstad for bringing this point to our attention.

drastic.³⁰ Similarly, ungrounded instances of (VD) are not in I_Ψ , and this restriction is justified as in the case of reflexivity.

5.1. Future work. We conclude by sketching some directions for further work. In the context of logical validity, the main open problem is the question of the global interpretability and, most importantly, of the \mathcal{L} -embedding of the theory of logical validity in the base theory. As for grounded consequence, the construction described by I_Ψ can be turned into an axiomatic theory. It would then be natural to study the relationships between this theory and the class of models extending $(\mathbb{N}, \mathsf{E}_\Psi, \mathsf{A}_\Psi)$. Finally, it would be interesting to relate irreflexive validity with paraconsistent theories of truth and validity.³¹

REFERENCES

- Beall JC and Julien Murzi (2013), “Two Flavours of Curry’s Paradox”, *The Journal of Philosophy* CX, 3, pp.143–65.
- Beklemishev, Lev (1995), “Iterated reflection versus iterated consistency”, *Annals of Pure and Applied Logic* 7, pp. 25–48.
- Beklemishev, Lev (2005), “Reflection principles and provability algebras in formal arithmetic”, *Russian Mathematical Surveys* 60, pp 197–268.
- Burgess, John P. (2014), “Friedman and the Axiomatization of Kripke’s Theory of Truth”, in N. Tennant (ed.), *Foundational Adventures. Essays in Honour of Harvey Friedman*, College Publications.
- Caret, Colin and Zach Weber (2015), “A Note on Contraction-Free Logic for Validity”, *Topoi* 34, pp. 63–74.
- Cobrerros, Pablo, Paul Egré, David Ripley, and Robert van Rooij (2014), “Reaching transparent truth”, *Mind*, 122, 488, pp. 841–866.
- Cook, Roy (2014), “There is No Paradox of Logical Validity”, *Logica Universalis*, published online 11 Jan 2014.
- Field, Hartry (2008), *Saving truth from paradox*, Oxford University Press, Oxford.
- Field, Hartry (forthcoming), “Disarming a Paradox of Validity”, forthcoming in the *Notre Dame Journal of Formal Logic*.
- French, Rohan (2016), “Structural Reflexivity and the Paradoxes of Self-Reference”, *Ergo* 3, 05.
- Greenough, Patrick (2001), “Free Assumptions and the Liar Paradox”, *American Philosophical Quarterly* 38, 2, pp. 115–135.
- Hájek, Petr and Pavel Pudlák (1998), *Metamathematics of First-Order Arithmetic*, Springer, Berlin.
- Halbach, Volker (2011), *Axiomatic Theories of Truth*, Cambridge University Press, Cambridge.
- Horsten, Leon (2011), *The Tarskian turn: deflationism and axiomatic truth*, MIT Press, Cambridge (MA).
- Ketland, Jeffrey (2012), “Validity as a primitive”, *Analysis* 72, 3, pp. 421–430.
- Kripke, Saul (1975), “Outline of a theory of truth”, *Journal of Philosophy* 72, 19, pp. 690–716.
- Mares, Edwin and Francesco Paoli (2014), “Logical Consequence and the Paradoxes”, *The Journal of Philosophical Logic* 43, 2, pp. 439–469.
- Leitgeb, Hannes (2005), “What Truth Depends On”, *Journal of Philosophical Logic* 34, pp. 155–192.
- Martin, Donald A. (2011), “Field’s *Saving Truth from Paradox*: Some Things It Doesn’t Do”, *Review of Symbolic Logic* 4, 3, pp. 339–347.
- Meadows, Toby (2014), “Fixed Points for Consequence Relations”, *Logique et Analyse* 57, 227, pp. 333–357.
- Montague, Richard (1963), “Syntactical treatments of modality, with corollaries on reflection principles and finite axiomatizability”, *Acta Philosophica Fennica* 16, pp. 153–67.

³⁰The standard counterpart of reflexivity in natural deduction is the rule of assumption (see [Negri and von Plato 2001], §1.3). If we accept a grounded picture of consequence, we can safely assume all the sentences of the base language, and apply all rules of inferences to them: the only sentences we cannot assume are the ungrounded ones, in line with our conception of consequence. Many thanks to Luca Tranchini for a useful discussion on this point.

³¹The latter research question has also been suggested in [French 2016].

- Moschovakis, Yiannis N. (1974), *Elementary Induction on Abstract Structures*, North-Holland and Elsevier, Amsterdam, London, New York.
- Murzi, Julien and Lionel Shapiro (2015), "Validity and Truth-Preservation", in T. Achourioti, K. Fujimoto, H. Galinon, and J. Martinez-Fernandez (eds.), *Unifying the Philosophy of Truth*, Springer.
- Negri, Sara and Jan von Plato (2001), *Structural Proof Theory*, Cambridge University Press, Cambridge.
- Nicolai, Carlo (forthcoming), "A dilemma for the reduction of semantic and modal notions", Unpublished Manuscript.
- Nicolai, Carlo (2016). "Equivalences for truth predicates", forthcoming in *The Review of Symbolic Logic*.
- Pailos, Federico and Diego Tajer (forthcoming), "Validity in a dialetheist framework", forthcoming in *Logique et Analyse*.
- Priest Graham and Richard Routley (1982), "Lessons from Pseudo Scotus", *Philosophical Studies* XLII, 2, pp. 189–99.
- Quine, W.V.O (1961). "Reply to Professor Marcus", *Synthese* 13, 4, pp. 323-330.
- Ripley, David (2012), "Conservatively Extending Classical Logic with Transparent Truth", *The Review of Symbolic Logic* 5, 2, pp. 354–378.
- Ripley, David (2015), "Comparing Substructural Theories of Truth", *Ergo* 2, pp. 299–328.
- Shapiro, Lionel (2011), "Deflating Logical Consequence", *The Philosophical Quarterly* LXI, 243, pp. 320–42.
- Schroeder-Heister, Peter (2012), "Paradoxes and Structural Rules", in C. Dutilh Novaes and O. T. Hjortland (eds.), *Insolubles and Consequences: Essays in honour of Stephen Read*, London, College Publications, pp. 203–211.
- Schroeder-Heister, Peter (2016), "Restricting Initial Sequents: The Trade-Offs Between Identity, Contraction and Cut", in R. Kahle, T. Strahm, and T. Studer (eds.), *Advances in Proof Theory* Birkhäuser 2016, pp. 339–351.
- Schwichtenberg, Helmut and Stanley S. Wainer (2012), *Proofs and Computation*, Cambridge University Press, Cambridge.
- Tennant, Neil (2015), "A new unified account of truth and paradox", *Mind*, 124 pp. 571–605.
- Visser, Albert (forthcoming), "The interpretability of inconsistency. Feferman's theorem and related results", Forthcoming in the *Bulletin of Symbolic Logic*.
- Yablo, Stephen (1982), "Grounding, Dependence, and Paradox", *Journal of Philosophical Logic* 11, pp. 117-137.
- Whittle, Bruno (2004), "Dialetheism, Logical Consequence and Hierarchy", *Analysis* LXI, pp. 320–42.
- Zardini, Elia (2011), "Truth without contra(d)iction" *Review of Symbolic Logic* 4, 4, pp. 498–535.
- Zardini, Elia (2013), "Naive *Modus Ponens*" *Journal of Philosophical Logic* 42 pp. 575–593.